

Data sheet

pColiExpress™ III Ligation Cloning & Expression Kit


Cat. No: BE011 (20 reactions)

Cat. No: BE012 (20 reactions)-Plus

Description

pColiExpress III is a DNA cloning vector for **periplasmic protein expression** in *E. coli*. All family pColiExpress DNA cloning vectors are based in a technology which allows efficiently cloning DNA fragments and the quick production of a large quantity of any desired protein.

pColiExpress III contains all elements required for periplasmic expression of **any desired protein** by OMPA Periplasmic Signal peptide and T7 RNA Polymerase Inducible System. pColiExpress III gives a His₆ tag at COOH end of the protein that allow the its purification. The vector has also an f1 origin of replication, an ampicillin resistance cassette, and a pBR322 origin of replication.



Unique restriction sites are showed.

Kit Components

Components	BE0011	BE0012 ⁽¹⁾
pColiExpress III (50 ng/μL) ⁽²⁾	20 μL	20 μL
5X T4 DNA Ligase Buffer	200 μL	200 μL
T4 DNA Ligase (5 U/μL)	20 μL	20 μL
Control Insert DNA (30 ng/μL)	10 μL	10 μL
pColiExpress III Control ⁽³⁾ (50 ng/μL)	5 μL	5 μL
Eco RV (10 U/μL)	-	20 μL
Hind III (10 U/μL)	-	20 μL
10X Buffer C	-	1.5 mL

(1) You need add at 5' end Hind III cleavage site in forward primer and Eco RV cleavage site at 5' end in reverse primer.

(2) Linearized DNA Vector.

(3) Circular vector. Empty DNA Vector.

pColiExpress vector family is a **ready to use** vector for a highly efficient cloning procedure. The vector is linearized, just for ligate with your PCR amplified with the recommending primers. Experimental background is less than 2%.

Features

T7 Promoter	4605-4621
Lac O	4627-4645
T7 transcription start	4621
Signal Peptide OmpA	4696-4777
His Tag	4786-4803
T7 terminator	4825-4872
Laclq	96-1175
Ori pBR322	2970-2356
Ampicillin resistance gene (ORF)	3133-3990
F1 Origin	4562-4115

(Continued on reverse side)

Distributed by:

Lab Unlimited
CARL STUART GROUP

Tallaght Business Park
Whitestown, Dublin 24,
Ireland
D24 RFK3

Tel: (01) 4523432
Fax: (01) 4523967

E-mail: info@labunlimited.com
Web: www.labunlimited.com

Quatro House, Frimley Road,
Camberley,
United Kingdom
GU16 7ER

Tel: 08452 30 40 30
Fax: 08452 30 50 30

E-mail: info@labunlimited.co.uk
Web: www.labunlimited.co.uk

canvax

Assay procedure

PCR restriction enzyme digestion

1. Your PCR reaction must be cleaned or loaded into agarose gel before.
2. Spin restriction enzymes and Buffer C to collect content at the bottom and set up the reaction as described below.

Match Reaction	Cloning Reaction
PCR cleaned	1 µg
10x Buffer C	7 µL
Eco RV (10 U/µL)	0.7 µL
Hind III (10 U/µL)	0.7 µL
Water (<i>Molecular Biology grade</i>)	up 70 µL

Cloning

3. Your PCR digestion must be cleaned and quantified before. Clean Easy PCR Purification kit (AN0063) is recommended.
4. Spin all kit components to collect content at the bottom of the tubes before set up the reaction.
5. Set up reaction as described below.

Match Reaction	Cloning Reaction	Control Reaction	Background Reaction
pColiExpress III vector (50 ng/µL)	1 µL	1 µL	1 µL
5x T4 DNA Ligase Buffer	2 µL	2 µL	2 µL
PCR Product*	X µL	-	-
Control Insert DNA	-	1 µL	-
T4 DNA ligase (5 Weiss units/µL)	1 µL	1 µL	1 µL
Water (<i>Molecular Biology grade</i>)	up 10 µL	up 10 µL	up 10 µL

*Relation vector: insert 1:5 is recommended.

6. Mix the reactions by pipetting.
7. Incubate one hour at Room Temperature (20-25°C).

Transformation

8. Centrifuge the tubes containing the reactions to collect content at the bottom of the tube. Add 15 µL of each reaction to a sterile 1.5 mL microcentrifuge tube on ice. Set up another tube on ice with 50 pg uncut plasmid for determination of the transformation efficiency of the competent cells (not supplied).
9. Place the competent cells in an ice bath until just thawed (about 10 to 15 minutes). Mix the cells by gently flicking the tube with your fingertips.
10. Carefully transfer 50 µL of cells into each tube prepared in Step 8.
11. Gently flick the tubes to mix and place them on ice for 30 minutes.
12. Heat-shock the cells for exactly 45 seconds in a water bath at exactly 42°C.
13. Immediately return the tubes to ice for 2 minutes and plate all transformation mix onto pre-warmed LB ampicillin plates. Incubate the plates overnight (12–16 hours) at 37°C.

Distributed by:

Lab Unlimited
CARL STUART GROUP

Tallaght Business Park
Whitestown, Dublin 24,
Ireland
D24 RFK3

Tel: (01) 4523432
Fax: (01) 4523967
E-mail: info@labunlimited.com
Web: www.labunlimited.com

Quatro House, Frimley Road,
Camberley,
United Kingdom
GU16 7ER

Tel: 08452 30 40 30
Fax: 08452 30 50 30
E-mail: info@labunlimited.co.uk
Web: www.labunlimited.co.uk